

Maryland Envirothon: Class Amphibia & Reptilia

KERRY WIXTED

WILDLIFE AND HERITAGE
SERVICE



March 2021

Amphibia Overview

- >40 species in Maryland
- Anura (frogs & toads)
- Caudata (salamanders & newts)
- Lay soft, jelly-like eggs (no shell)
- Have larval state with gills
- Breathe & drink through skin



Gray treefrog by Kerry Wixted

Note: This guide is an overview of select species found in Maryland.

Anura

- ~20 species in Maryland
- Frogs & toads
- Short-bodied & tailless (as adults)
- Typically lay eggs in water & hatch into aquatic larvae



Green treefrog by Kerry Wixted

Family Bufonidae (Toads)



Photo by Kerry Wixted

American Toad (*Anaxyrus americanus*)

2-3.5"; typically 1-2 spots/ wart; parotoid gland is separated from the cranial crest or connected narrowly by a spur; enlarged warts on tibia; **Call:** an elongated trill or whir lasting 5-30 seconds and resembles a simultaneous whistle and hum



Photo by Judy Gallagher CC by 2.0

Fowler's Toad (*Anaxyrus fowleri*)

2-3"; typically 3+ spots/ wart; parotoid gland is in contact w/ the cranial crest; **Call:** a short, brash and whiny call lasting 2-4 seconds

Family Hylidae (Treefrogs)



Spring Peeper

(*Pseudacris crucifer*)

0.75 - 1.25"; Brown, tan, or yellowish with dark X-shaped mark on back; Dark bar between eye; Mask from nose through eye and tympanum, often extending down side

Call: Clear, shrill, high-pitched whistle or peep



Gray Treefrog & Cope's Gray Treefrog

(*Hyla versicolor* & *Hyla chrysoscelis*)

1.25 - 2" (Identical in appearance); Gray to white with darker streaking, resembling a tree knot; Cream square below each eye; Inner thigh yellow or orange; enlarged toe pads; **Call** (*H. versicolor*): Musical, bird-like trill **Call** (*H. chrysoscelis*): Rapid, more ratchet-like trill

Family Ranidae (True Frogs)



Northern Green Frog (*Lithobates clamitans*)

2.25 - 3.5"; Underside white (or yellow in breeding males); dorsolateral ridges on the back end before reaching the groin **Call:** Short and accented multi-part call, bursts of sound akin to the plucking of a banjo string



Bullfrog (*Lithobates catesbeianus*)

3.5 - 6+"; Underside white with mottling (or yellow in breeding males); no dorsolateral ridge; **Call:** Deep pitched, resembling the bellow of a bull

Caudata

- > 20 species in Maryland
- North America has more salamander species than the rest of the world combined!
- Undergo metamorphosis



Marbled salamander by Kerry Wixted

Family Ambystomidae (Mole Salamanders)



Marbled Salamander (*Ambystoma maculatum*)

3.5 – 5.5”; Has a stout body with light gray (female) or white (male) crossbars on dark gray to black body; breeds in the fall at the edge of small, shallow wetlands (vernal pools); lives in woodland habitat and spends most time underground

Ambystomidae (Mole Salamanders)

By Andrew Hoffman CC by NC ND2.0



Eastern Tiger Salamander (*Ambystoma tigrinum*)

7 – 8.25"; Dull black to deep brown with irregular light olive or yellowish brown spots extending down sides; breeds in winter; **State Endangered**



Spotted Salamander (*Ambystoma maculatum*)

4.5 - 8"; Black or dark gray with paired yellow spots; spots occasionally orange or absent; found in hardwood forests & breeds in early spring

Family Cryptobranchidae

Eastern Hellbender

(*Cryptobranchus alleganiensis*)

- State Endangered
- Up to 29" in length
- Flat head, sides w/ wrinkled flesh folds
- Clear, flowing streams



Photos By Dave Kazyak

Family Plethodontidae

Eastern Red-backed Salamander (*Plethodon cinereus*)

- 2.25 - 4" in length
- Salt & pepper pattern on belly
- Has striped (with reddish mid-dorsal stripe) & leadback variants (uniformly dark gray-black)
- One of most abundant vertebrates in North America



Family Salamandridae

Red-spotted Newt

(*Notophthalmus viridescens viridescens*)

- 2.25 – 4.8” in length
- Aquatic or semi-aquatic adults olive green with red spots outlined in black
- Terrestrial juveniles (efts) orange-red or brownish with red spots outlined in black
- Skin is granular in appearance compared to other salamander species



Reptilia Overview

- >50 species in Maryland
- Testudines (turtles)
- Squamata (lizards & snakes)
- Lack aquatic larval stage
- Have scaly skin
- Lay eggs w/ shells or give live birth



Eastern Snapping turtle by Kerry Wixted

Note: This guide is an overview of select species found in Maryland.

Testudines

- ~22 species in Maryland
- Upper part of shell= carapace
- Lower part of shell= plastron
- Scales= scutes
- Largest species= leatherback sea turtle



Family: Chelydridae

Eastern Snapping Turtle (*Chelydra serpentina*)

- 8-19" carapace length
- Long neck and big head with powerful jaws
- Skin covered in many bumps and tubercles
- Long tail with three rows of spiky tubercles
- Rear of shell is saw-toothed



Juvenile & adult snapping turtle by Kerry Wixted

Family Emydidae

- ~12 species in Maryland
- Includes box turtle, sliders, terrapins, and more
- Many are adapted for swimming



Eastern painted turtles and northern red-bellied cooters on log



By Kerry Wixted

Pond Turtles

By Matthew Beziat CC by 2.0



Red-eared Slider

(*Trachemys scripta elegans*)

5-8"; **invasive**; plastron yellow with dark blotches; has broad reddish-stripe behind eye in most individuals



Eastern Painted Turtle

(*Chrysemys picta picta*)

4.5-8"; plastron yellow and lightly spotted; scutes on carapace almost in straight lines across back

Spotted Turtle (*Clemmys guttata*)



By Kerry Wixted



- 3.5-4.5" carapace
- Yellow spots on shell

- Head & neck w/ yellow or orange spots
- Found in shallow, marshy meadows and bogs, swamps and small ponds

Bog Turtle (*Glyptemys muhlenbergii*)

- Federally Threatened
- State Threatened
- 3-4" carapace
- Orange patches on sides of head
- Lives in sphagnum bogs and open swamps



By USFWS

Eastern Box Turtle (*Terrapene carolina*)

- 4.5 - 6" carapace
- High, domelike shell
- Extremely variable in color & pattern
- 4 toes on hindlimbs
- Female:
 - Eyes normally brown
 - Flat plastron
- Male:
 - Eyes often red
 - Central concave on plastron
- Woodlands and thickets= habitat



Northern Diamond-backed Terrapin (*Malaclemmys terrapin terrapin*)

- Female: 6 - 9.3" carapace
- Male: 4 - 5.5" carapace
- Concentric grooves and ridges or concentric dark and light markings on each carapace scute
- Skin coloration is cream or light silver to dark gray or black, often with dark spotting
- Light-colored beak, sometimes with dark "mustache"
- MD state reptile
- Coastal marsh habitat



[This Photo](#) by Unknown Author is licensed under [CC BY-SA](#)

Squamata

- ~33 species in Maryland
- “Scaled reptiles”
- Lizards & snakes
- Introduced=
 - Mediterranean Gecko



M. Gecko by Jim Brighton, Maryland Biodiversity Project



Family Phrynosomatidae

Eastern Fence Lizard (*Sceloporus undulatus*)

- 4-7.25" in length
- Heavily keeled scales
- Dark, lateral cross bars (horizontal stripes)
- Females are gray with darker cross bars
- Males may be brownish with very faint crossbars (if present) and an iridescent blue and black patch at the base of the throat and on each side of the belly



Male Eastern Fence Lizard by Kerry Wixted

Family Teiidae

Six-lined Racerunner

(*Aspidoscelis sexlineatus*)


- 6-10.5" in length
- Slender body w/ very long tail
- Six light-colored stripes from the head through the tail
- Sandpaper-like scales on body; keeled scales on tail which appear rough
- Open areas with sand or loose soil= habitat



Six-lined Racerunner by Kerry Wixted

Snakes of Maryland

Two species, the Eastern Copperhead and the Timber Rattlesnake, are pit vipers (Subfamily Crotalinae) and possess medically significant venom. Pit vipers are named for the heat-seeking pit between each eye and nostril. Both of Maryland's pit vipers should be treated with caution and should not be approached or handled. The graphic on the left, and the associated table below, show characteristics of pit vipers in Maryland versus other Maryland snakes. These characteristics do not work for venomous snakes outside of Maryland. For more information on snakes, please visit: dnr.maryland.gov/wildlife



Other Maryland Snake ID Characters	Pit Viper ID Characters
1. Round pupils	1. Elliptical pupils
2. No sensory pit	2. Sensory pit between eye and nostril
3. Head slightly wider than neck*	3. Head much wider than neck*
4. Divided anal plate scale**	4. Single anal plate scale**
5. Double row of scales on the underside of tail **	5. Single scales on underside of tail**

* Some snake species, like Eastern Ratsnakes, Common Watersnakes, and Common Gartersnakes will flatten their heads when threatened to mimic pit vipers.

** These characteristics are helpful when examining shed snake skins.

Family Colubridae

- Egg-laying snakes
- Includes ratsnakes, racers, green snakes, and more



Eastern Ratsnake By Kerry Wixted

Eastern Ratsnake (*Pantherophis alleghaniensis*)

Photo by Robert Warren



Photo by Richard Orr



Juvenile: Grayish with dark blotches along back. Has dark bar between eyes that continues past the eye to corner of mouth.

Adult: 42-72"; Uniform black on top or with faint traces of a spotted pattern. Throat and underside light colored. Adept climber.

Northern Black Racer (*Coluber constrictor constrictor*)

Photo by Bill Hubick, MD Biodiversity Project



Photo by Lisa Reid CC by NC ND 2.0



Juvenile: Strongly patterned, like watersnakes and ratsnakes, but have more blotches down the back, a set of blotches along the centerline and proportionally larger eyes, an adaptation for open field hunting.


Adult: 36 - 60"; Sleek in appearance. Plain black above and steely gray to black below. Often white coloration seen on the chin. Large prominent eye.

Family Dipsadidae

- Formerly in Colubridae
- Rear-fanged snakes
- Small to moderately sized

Eastern Wormsnake
(*Carphophis amoenus amoenus*)

7.5-11"; brown above w/ pink belly; tail ends in sharp spine; scales smooth; found in woodland habitats



By Kerry Wixted

Family Dipsadidae: Ring-necked Snakes

Northern Ring-necked Snake

(*Diadophis punctatus edwardsii*)

10-15"; Back is a uniform dark gray, bluish black or brown. Has cream-to-yellow or orange neck collar and yellow belly, which may grade to red towards the tail. Has complete neck ring and belly has no or small black spots down the midline.

Southern Ring-necked Snake

(*Diadophis punctatus punctatus*)

10-15"; Back is a uniform dark gray, bluish black or brown. Has cream-to-yellow or orange neck collar and yellow belly, which may grade to red towards the tail. Has broken or notched neck ring behind head and belly has large, black half-moon shaped spots down the midline.



Photo by: Linh Phu



Photo by: Matt Close

Eastern Hog-nosed Snake (*Heterodon platirhinos*)

- 20-30" in length
- Coloration variable with yellow, brown, gray, orange or red on a dark splotch-patterned back
- Some are an un-patterned ebony black or dark grey
- Snout upturned
- Often will flatten out head, hiss, pretend to strike and/or play dead if disturbed



Photo by Andrew Hoffman CC by NC ND 2.0

Family Natricidae

- Formerly in Colubridae
- Harmless, live-bearing snakes

DeKay's Brownsnake (*Storeria dekayi*)

Small, brownish/gray snake with small, paired black spots running down the back. A thin dark line extends from the back of the head behind the eye to the base of the jaw. Juveniles have a light band across their neck.




Illustration showing the presence of an incomplete cream-colored collar on the juvenile DeKay's Brownsnake.



Photo by: Mark Nenadov CC by 2.0

Northern Watersnake (*Nerodia sipedon sipedon*)

Photo by Andrew DuBois CC by NC 2.0



Photo by Judy Gallagher CC by 2.0



22-42"; pattern highly variable with variations including bands that are thicker at the top and thinner at the bottom, incomplete bands or no bands. Has dark lip stripes. Often mistaken for cottonmouth (water moccasin) which does not occur in Maryland.

Family Crotalinae

- All are venomous
- Heads are wider than the neck
- Possess facial pits between eye & nostrils
- Scales under tail are in a single row



Timber Rattlesnake by Kerry Wixted

Eastern Copperhead (*Agkistrodon contortrix*)

- 24-36"
- Has Hershey-kiss shaped markings that are skinny at the top and wider at the base
- Head is coppery in color with a contrasting upper lip
- Juveniles have bright yellow caudal lure



Timber Rattlesnake (*Crotalus horridus*)


- 36-60"
- Found in western Maryland
- Color variable from light and striped to almost dark black with jagged stripes
- Has distinct rattle at end of tail
- Found in wooded areas near rocky outcroppings



Photo by: Peter Paplanus CC by 2.0

Juvenile Snake Comparisons

Photo by: Andrew Hoffman CC by NC ND 2.0



Andrew Hoffman, 2008

Northern Copperhead




North American Racer

Photo by: Bill Hubick, Maryland Biodiversity Project

Photo by: Robert Warren



Eastern Ratsnake



Common Watersnake

Photo by: Andrew DuBois CC by NC 2.0

Preventing Snake Bites

Snake bites are not common, but it is important to be aware they can occur and how to deal with them. Below are some tips on how to avoid being bitten:

1. Learn how to identify the copperhead and the timber rattlesnake.
2. When hiking or camping, watch where you put your hands and feet. Be mindful of where you sit and where you place your sleeping bag.
3. Wear suitable clothing when hiking, especially through tall grass or heavy brush. Long pants and heavy boots are usually best in tall grass and heavy brush.
4. Avoid rock piles and stacks of old boards or wood in forested areas as snakes use these areas frequently, especially sunny areas with canopy gaps.
5. Be careful working around brush piles or other debris. Use a rake or long handled tool to move brush, debris, or other material before picking it up.
6. Never handle venomous snakes, alive or dead.
7. Leave snakes alone. Many bites occur when people attempt to capture or kill venomous snakes.

If bitten by a venomous snake, immediately contact the Maryland Poison Center at 800-222-1222. The Maryland Poison Center is open 24 hours a day and is staffed by pharmacists or nurses who are specially trained and certified in emergency poisoning and overdose cases. Do not apply ice, a tourniquet, or make an incision around the wound as these methods do not work and may cause complications.



Ranavirus

- Viruses that impact cold-blooded animals
- Larval wood frogs, spotted salamanders and marbled salamanders have experienced the most mortality
- 1st significant box turtle die-off in Maryland in 2008
- Disinfection of equipment (boots, waders, etc) before entering wetlands is recommended
 - Protocol: <http://northeastparc.org/disinfection-protocol/>



Eastern box turtle w/ Ranavirus by
Scott Farnsworth

For Additional Information


Special thanks to Rachel Gauza for assistance putting together this guide. For more information on Maryland's reptiles and amphibians, please visit our webpage at: dnr.maryland.gov/wildlife Additional information can be found in the following resources.

Book Resources:

- The Maryland Amphibian and Reptile Atlas
- Peterson Field Guide to Reptiles and Amphibians of Eastern and Central North America, 4th Ed

Online Resources:

- [Maryland Envirothon \(Wildlife and Heritage Service\)](#)
- [Maryland Biodiversity Project \(Amphibians\)](#)
- [Maryland Biodiversity Project \(Reptiles\)](#)
- [Virginia Herpetological Society](#)



Larry Hogan, Governor Jeannie Haddaway-Riccio, Secretary
Wildlife and Heritage Service * 580 Taylor Ave, E-1* Annapolis, MD 21401
410-260-8540
dnr.maryland.gov/wildlife