




Dissolved oxygen conditions in the Maryland Chesapeake Bay mainstem were average in early August. The hypoxic water volume (areas below 2 mg/l oxygen) was approximately 1.15 cubic miles, which is close to the early August 1985-2015 average of 1.13 cubic miles. No anoxic zones (areas below 0.2 mg/l) were detected, and none have been observed since the summer of 2014. Hypoxia abated from Virginia waters and constricted laterally along the bay's channel versus the late July results. Salinity levels are higher this year compared to last, indicating lower rainfall, and less nutrient runoff which may exacerbate hypoxia.

In the beginning of June, National Oceanic and Atmospheric Administration, United States Geological Survey, University of Maryland Center for Environmental Science and University of Michigan scientists predicted a close to average sized hypoxic volume for the bay due to lower spring flows (January-May) and nitrogen loading from the Susquehanna and Potomac rivers.



Late July



Early August

For more information:

- Maryland Department of Natural Resources - Our Waters Page. dnr.maryland.gov/waters
- Press Release for the 2016 Chesapeake Bay Hypoxic Zone Forecast. 1.usa.gov/28QHmEc

The facilities and services of the Maryland Department of Natural Resources are available to all without regard to race, color, religion, sex, sexual orientation, age, national origin or physical or mental disability.

This document is available in alternative format upon request from a qualified individual with disability.

Crabs, fish, oysters and other creatures in the Chesapeake Bay require oxygen to survive. Scientists and natural resource managers study the volume and duration of bay hypoxia to determine possible impacts to bay life.

Each year (June-September), the Maryland Department of Natural Resources computes these volumes from data collected by Maryland and Virginia monitoring teams. Data collection is funded by these states and the Environmental Protection Agency's Chesapeake Bay Program. Bay hypoxia monitoring and reporting will continue through September.

Posted: August 17, 2016

